



IX Congresso Nazionale AIPP

**Mind the gap: l'intervento precoce
tra continuità evolutiva, discontinuità
diagnostiche e multiculturalità.**

Bari, 27-28-29 Settembre 2023
Università degli Studi di Bari "Aldo Moro"

DISCLOSURE INFORMATION

Marina Mancini

Dichiaro che negli ultimi due anni non ho avuto rapporti di finanziamento con soggetti portatori di interessi commerciali in campo sanitario



Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP


Dipartimento
Salute Mentale



LA NECESSITÀ DI INTERVENTI EVIDENCE-BASED SULL' ABUSO DI SOSTANZE NEI GIOVANI A RISCHIO E ALL'ESORDIO PSICOPATOLOGICO: L'ESPERIENZA CLINICA NEL CENTRO DEDICATO GIPPSI

Dott.ssa Marina Mancini, Dirigente Psicologo UOSVD Esordi Psicotici, ASL Bari

INTRODUZIONE





Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP

Dipartimento
Salute Mentale



FATTORI STRESSANTI/AMBIENTALI

- 1) Separazione dei genitori
- 2) Uso di sostanze



Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP

Dipartimento
Salute Mentale



Psychiatry

RESEARCH ARTICLE

Open Access

Common stressful life events and difficulties are associated with mental health symptoms and substance use in young adolescents

Nancy CP Low^{1*}, Erika Dugas², Erin O'Loughlin², Daniel Rodriguez³, Gisele Contro^{2,4}, Michael Chouinard⁵
and Jennifer O'Loughlin^{2,4,6}

Schizophrenia Bulletin vol. 42 no. 5 pp. 1262–1269, 2016
doi:10.1093/schbul/sbw003
Advance Access publication February 15, 2016

Meta-analysis of the Association Between the Level of Cannabis Use and Risk of Psychosis

Arianna Marconi¹, Marta Di Forti¹, Cathryn M. Lewis², Robin M. Murray¹, and Evangelos Vassos^{*,2}

¹Department of Psychosis Studies, King's College London, Institute of Psychiatry Psychology & Neuroscience, London, UK; ²King's College London, Institute of Psychiatry Psychology & Neuroscience, MRC SGDP Centre, London, UK

*To whom correspondence should be addressed; King's College London, Institute of Psychiatry Psychology & Neuroscience, MRC SGDP Centre, Box P082, De Crespigny Park, London SE5 8AF, UK; tel: +44-20-7848-5433, fax: +44-20-7848-0866, e-mail: evangelos.vassos@kcl.ac.uk



INTRODUZIONE

- Diversi studi hanno evidenziato come nei soggetti con disturbo psicotico l'uso di sostanze sia significativamente più elevato rispetto alla popolazione generale. (McCraedie RG., 2002; Tomassini A., 2012)
- Inoltre, è stata osservata un'alta prevalenza di uso di sostanze, in particolare cannabis e alcool, tra gli individui con un primo episodio psicotico. (Verdoux H., 2005; Tomassini A., 2012)



INTRODUZIONE

- L'uso di sostanze è stato associato con una più giovane età all'esordio, genere maschile, mancanza di deficit cognitivi, ricaduta precoce e maggior gravità dei sintomi.
- L'aumentato uso di cannabis ha anche un impatto sul decorso a lungo termine del disturbo, in termini di mancata remissione e gravità dei sintomi.
- Il 52% dei soggetti valutati in uno studio del 2007 mostrava una comorbidità con un disturbo da uso di sostanze, nel 32% dei casi la sostanza era la cannabis e nel 35% l'alcool (Addington & Addington).



Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP

Dipartimento
Salute Mentale



INTRODUZIONE

European Psychiatry

www.cambridge.org/epa

Original Article

Cite this article: Ayerbe L, Pérez-Piñar M, Foguet-Boreu Q, Ayís S (2020). Psychosis in children of separated parents: A systematic review and meta-analysis. *European Psychiatry*, 63(1), e3, 1–9
<https://doi.org/10.1192/j.eurpsy.2019.15>

Received: 29 October 2019

Revised: 26 November 2019

Accepted: 28 November 2019

Key words:

childhood adversities; divorce; parental separation; psychotic disorders; schizophrenia

Author for correspondence:

Luis Ayerbe,

E-mail: lgarcia-morzon@qmul.ac.uk

Psychosis in children of separated parents: A systematic review and meta-analysis

Luis Ayerbe^{1,2,3}, María Pérez-Piñar³, Quintí Foguet-Boreu^{4,5} and Salma Ayís^{6,7,8}

¹Department of Public Health and Primary Care, University of Cambridge, Cambridge, United Kingdom; ²Centre of Primary Care and Public Health, Queen Mary University of London, London, United Kingdom; ³Carnarvon Medical Centre, Southend-on-Sea, United Kingdom; ⁴Department of Psychiatry, Vic University Hospital, Vic, Spain; ⁵Faculty of Health Sciences and Welfare, University of Vic-Central University of Catalonia (UVic-UCC), Vic, Spain; ⁶School of Population Health and Environmental Sciences, King's College London, London, United Kingdom; ⁷National Institute for Health Research Biomedical Research Centre, Guy's and St Thomas' NHS Foundation Trust, London, United Kingdom and ⁸National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care, South London at King's College Hospital NHS Foundation Trust, London, United Kingdom

Abstract

Background. Parental separation is a very common childhood adversity. The association between other adverse childhood experiences and an increased risk of psychosis has been reported. However, the evidence on the risk of psychosis for children of separated parents is limited. In this systematic review, cohort, case-control, and cross-sectional studies, comparing the risk of psychotic disorders for people with and without separated parents, were searched, critically appraised, and summarized.

Methods. Studies were searched in PubMed, EMBASE, PsycINFO, and the Web of Science, from database inception to September 2019. A meta-analysis, using random-effects models, was undertaken to obtain pooled estimates of the risk of psychosis among participants with separated parents.

Results. Twelve studies, with 305,652 participants from 22 countries, were included in the review. A significantly increased risk of psychosis for those with separated parents was observed, with a pooled odds ratio: 1.53 (95% confidence interval [CI]: 1.29–1.76), $p < 0.001$. The association remained significant when cohort, case-control, and cross-sectional studies were analyzed separately. The five cohort studies included in this review showed an increased risk of psychosis with odds ratio: 1.47 (95% CI: 1.26–1.69), $p < 0.001$.


Conclusions. Parental separation is a common childhood adversity associated with an increased risk of psychosis. Although the risk for an individual child of separated parents is still low, given the high proportion of couples that separate, the increased rates of psychosis may be substantial in the population. Further studies on the risk of psychosis in those with separated parents, and the

Diversi studi hanno evidenziato l'associazione tra l'evento Separazione o divorzio dei genitori e l'abuso di sostanze nella giovane età, oltre all'aumento del rischio di psicosi (Sands et al, 2017; Schlossarek et al, 2016).

Dott.ssa Marina Mancini, Dirigente Psicologo UOSVD Esordi Psicotici, ASL Bari




L'ESPERIENZA DEL CENTRO GIPPSI





L'ESPERIENZA DEL CENTRO GIPPSI

APPROFONDIMENTO PSICODIAGNOSTICO



L'ESPERIENZA DEL CENTRO GIPPSI

N= 112 (52 UHR, 60 ESORDI)

Età media= 21±3.5




Figura 1. Interazione tra abuso di alcool ed evento «separazione dei genitori» sul Global Severity Index del test SCL-90 che rappresenta un indicatore globale dell'intensità del disagio psichico lamentato dal rispondente




Figura 2. Interazione tra abuso di alcool ed evento «separazione dei genitori» sul Global Severity Index del test SCL-90 nel sottogruppo Esordi

L'ESPERIENZA DEL CENTRO GIPPSI




Figura 3. Interazione tra abuso di alcool ed evento «separazione dei genitori» sul Positive Symptom Total del SCL-90 che indica il numero totale di sintomi riportati dal rispondente





Figura 4. Interazione tra abuso di alcool ed evento «separazione dei genitori» sul Positive Symptom Total del SCL-90 che indica il numero totale di sintomi nel sottogruppo Esordi

L'ESPERIENZA DEL CENTRO GIPPSI





Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP

Dipartimento
Salute Mentale



QUALI POSSIBILI INTERVENTI?

PSICOTERAPIA
INDIVIDUALE

INTERVENTI SULLE
FAMIGLIE

PSICOEDUCAZIONE
SULL'USO DI
SOSTANZE



Fondazione
E.P.A.S.S.S.



IX Congresso Nazionale AIPP

Dipartimento
Salute Mentale



GRAZIE PER L'ATTENZIONE!

Dott.ssa Marina Mancini, Dirigente Psicologo UOSVD Esordi Psicotici, ASL Bari